


Table I. Chiral Catalysis in Additions of Alkyl- or Aryllithiums to Aldehydes in 150 mL of Et₂O at -120 °C

run no	diamine		RLi		R'CHO		C ₆ H ₅ CH(OH)R				$\Delta(\Delta G^\ddagger)^c$ kcal mol ⁻¹
	struct	mmol	R	mmol	R'	mmol	% optical yield (ee)	yield, %	config		
1	(R,R)-1	5	CH ₃ (CH ₂) ₃	4.4	C ₆ H ₅	1.2	95	89	73	R	1.0
2	(R,R)-1	3.8	C ₆ H ₅	3.1	CH ₃ (CH ₂) ₃	1.2	43	55	47	S	0.3
3	(R,R)-1	3.6	CH ₃ (CH ₂) ₂	3.4	C ₆ H ₅	1.2	59	63	63	R	0.4
4	(R,R)-1	3.4	CH ₃ CH ₂	3.3	C ₆ H ₅	1.2	66	67	75	R	0.5
5	(R,R)-1	3.9	CH ₃	3.1	C ₆ H ₅	1.2	36	35	35	R	0.2
6 ^d	(R)-2	3.3	CH ₃ (CH ₂) ₃	3.0	C ₆ H ₅	1.0	58	55	71	R	0.4
7	(R)-2	3.4	CH ₃ (CH ₂) ₂	3.0	C ₆ H ₅	1.2	53	54	75	R	0.4
8	(R)-2	3.3	CH ₃ CH ₂	2.2	C ₆ H ₅	1.2	30	30	73	R	0.2
9	(R)-2	3.6	CH ₃	3.0	C ₆ H ₅	1.2	23	22	65	R	0.1

^a Reference 8 lists maximum rotations used. ^b From ¹H NMR of (+)- α -methoxy- α -[(trifluoromethyl)phenyl]acetic esters. ^c $\Delta(\Delta G^\ddagger) = RT \ln$ [dominant enantiomer/subordinate enantiomer]. Values of ee determined by the two methods were averaged. ^d Volume of Et₂O was 20 mL.

than pentanal as the aldehyde component.

Without exception, the configurational bias is correlated by the formulations of generalized transition states 4 and 5 to give 6.




Although we examined only 14 out of many possible combinations of solvents, temperatures, reagents, catalysts, and orders of attaching groups to the asymmetric centers, the fact that all 14 reactions fit this correlation suggests that a single mechanistic feature dominates the chiral tilt. Examination of CPK molecular models of 4 with R = CH₃(CH₂)₃ and R' = C₆H₅ in advance of experiment provided the correct prediction of the configuration of the dominant product. Because of the rigid polyspirane structures of the presumed transition states, the only degree of freedom is the placement of the H and R' groups. In 4 and 5, that diastereomer is formulated in which the H is directed toward the face of the upper left naphthalene and, in 4, R is directed alongside the face of the lower right naphthalene. Models of 4 and 5 appear less sterically constrained than those of their diastereomers in which the positions of the H and R' groups are interchanged. Since N: rapidly inverts, catalyst (R,R)-1, in effect, contains three C₂ axes, and 4 can be generated four different ways from (R,R)-1, R'CHO, and RLi. Since catalyst (R)-2 in effect contains one C₂ axis, 5 can be generated two different ways from (R)-2, R'CHO, and RLi. This degeneracy reduces the number of transition states that compete with one another, the averaging of compensating effects, and probably is at least partly responsible for the high chiral recognition of the catalysts.

This correlation shows the following similarities to the rule of steric control of asymmetric induction for additions to carbonyl groups adjacent to chiral centers.¹² In both correlations, the differences in free energies between the diastereomeric transition states cover about the same range and depend on steric effects. An inversion in the order of attachment of groups to the forming asymmetric center inverts the configuration of that center.

Seebach et al. have used optically active amino ethers derived from tartaric acid to get 56–15% ee in stoichiometric catalysis of additions of organometallics to aldehydes.¹³ However, Mukaiyama et al. reported the highest observed stoichiometric catalysis of additions of alkylorganometallics to aldehydes to provide

alcohols.¹⁴ When RLi complexed by (S,S)-7 was added to C₆H₅CHO, the optical yields varied from 83¹⁵ to 21% ee, and the



dominant isomer produced depended on the R of RLi and the solvent. The highest optical yield (92% ee) involved Et₂Mg reacting with C₆H₅CHO.

(14) Mukaiyama, T.; Soai, K.; Sato, T.; Shimizu, H.; Suzuki, K. *J. Am. Chem. Soc.* 1979, 101, 1455-1460.

(15) These authors based their optical yields on a maximum rotation for C₆H₅CH(OH)(CH₂)₃CH₃ of $[\alpha]_D^{25} 31.3^\circ$ (c 3, C₆H₆). We have corrected their ee values to a maximum of $[\alpha]_D^{25} 35.7^\circ$ (c 3, C₆H₆) so that their results and ours can be compared (see ref 9).


Aromatic Polycyclic Benzenoid Tetraanions: Pyrene and Perylene Anions Revisited

Abraham Minsky, Joseph Klein, and Mordecai Rabinovitz*

Department of Organic Chemistry
The Hebrew University of Jerusalem
Jerusalem 91904, Israel
Received February 17, 1981

Recently, it has been reported that four-electron reduction of conjugated hydrocarbons is feasible.¹ Conjugated systems of (4n + 2) π electrons in their periphery are of particular interest, being predicted to acquire aromatic character as a consequence of a four-electron reduction process.²

No information exists on quadruply charged ions which originate from 4n + 2 peripheries of the "ultimate aromatics", i.e., benzenoid systems. We wish to report a reduction of two polycyclic benzenoid (4n + 2) π systems which occupy a cardinal position in the understanding of aromaticity, i.e., pyrene (1) and perylene (2) leading to the formation of the respective tetraanions. These



(1) (a) Paquette, L. A.; Ewing, G. D.; Traynor, S.; Hardlik, J. M. *J. Am. Chem. Soc.* 1977, 99, 6115-6117. (b) Huber, H.; Müllen, K.; Wennerström, O. *Angew. Chem., Int. Ed. Engl.* 1980, 19, 624-625. (c) Müllen, K.; Oth, J. M.; Engels, H. W.; Vogel, E. *Ibid.* 1979, 18, 229-231. (d) Becker, B. Ch.; Huber, W.; Müllen, K. *J. Am. Chem. Soc.* 1980, 102, 7803-7805. (e) Huber, W.; Müllen, K. *J. Chem. Soc., Chem. Commun.* 1980, 698-700.

(2) Platt, J. R. *J. Chem. Phys.* 1954, 22, 1448-1452.

(12) Cram, D. J.; Abd Elhafez, F. A. *J. Am. Chem. Soc.* 1952, 74, 5828-5835.

(13) Seebach, D.; Crass, G.; Wilka, E. M.; Hilvert, D.; Brunner, E. *Helv. Chim. Acta* 1979, 62, 2695-2698.

Table I. ¹H NMR Chemical Shifts of Pyrene (1), Perylene (2), and Tetraanions 3 and 4⁷

	pyrene (1)	pyrene ^a dianion	pyrene tetraanion (3)	perylene (2)	perylene tetraanion (4)	isopyrene ^a	isopyrene dianion ^a
center of gravity of ¹ H NMR bands	8.15	0.97	5.3	7.89	5.35	8.05	6.72
Δ(δ) from the neutral compd		7.18	2.85		2.54		1.33 ^b

^a Data taken from ref 3. ^b The isopyrene has a LUMO with a nodal plane which does not bisect the central carbon atoms: so this orbital is not a pure peripheral one, a fact which explains the reduced paratropicity revealed in isopyrene dianion.³

two systems have already been subjected in the past to reduction by metals. A doubly charged paratropic species was detected as a dilithium salt³ of **1** to yield **1**²⁻ and a disodium salt was reported to be formed by metal reduction of **2**.⁴ Comparison of the reported ¹H NMR spectra of these two charged systems shows a striking difference between them. Upon reduction of **1** to **1**²⁻ the proton chemical shift's center of gravity Δ(δ) shifts upfield by 7.18 ppm, while the total change in reduction of **2** is only 2.54 ppm. This difference cannot be ascribed to the nonsimilarity of the π periphery of these hydrocarbons as they accommodate 14- and 18π electrons and **1** and **2**, respectively.⁵ Simple HMO of **1** and **2** show that each one of them has a LUMO orbital exhibiting a nodal plane through the central carbon atoms.⁶ This leads to an important conclusion: in **1**²⁻ and **2**²⁻ the molecular orbitals which are occupied by the two additional electrons represent those of a pure peripheral annulene.

This discrepancy tempted us to study the metal reduction of **1** and **2**. Reduction of **1** with sodium metal at room temperature (THF-*d*₈, sealed degassed tube) was monitored by ¹H NMR spectroscopy.⁷ After 4 days the deep purple solution exhibited a well-resolved AB₂M spectrum; the AB₂ part is centered at 5.68 ppm, the M part at 4.40 ppm, and the center of gravity of the entire spectrum at 5.30 ppm (Table I). This spectrum remains unchanged for weeks. During the first 4 days no proton bands were detected, presumably due to the formation of radical species and fast bimolecular electron-transfer processes. This ¹H NMR spectrum is entirely different from the spectrum of the paratropic species **1**²⁻ reported previously which was obtained with lithium metal at -80 °C.⁸ An experiment conducted in dimethoxyethane showed the same spectrum. Oxidation of the anion solutions with O₂ afforded **1** as a sole product. These experiments indicate that the ¹H NMR bands could not originate from a reaction product of the anion itself or the anion and solvent. Quenching with D₂O afforded tetradeuteriotetrahydropyrene (*m/e* 210, P + 1 = 211) as shown by electron capture and chemical ionization mass spectrometry,⁹ accompanied by a mixture of partially dehydrogenated products (*m/e* 202-209). We followed the formation of the various species in the reaction of pyrene and sodium by repeated sampling experiments. Samples were taken from the reaction mixture, quenched with water, and titrated potentiometrically to follow the formation of base. The amount of sodium was followed by atomic absorption. The amount of base and sodium increased with time, and after 4 days 3.3-4.0 equiv of base and 3.2-4.0 atoms of sodium were detected per molecule of **1**. No further increase of base and sodium could be detected even after 14 days. These results clearly indicate the formation of a quadruply charged anion **3**, i.e., **1**⁴⁻.¹⁰

Similar experiments were conducted with the perylene system (**2**). The ¹H NMR spectrum⁷ of the purple solution formed by contact between **2** and sodium after 1 week afforded an identical ¹H NMR spectrum to that reported by Lawler et al.⁴ (Table I) and assigned to the formation of a dianion. Quenching with D₂O as mentioned before⁹ showed clearly the formation of a tetradeuteriotetrahydroperylene species (*m/e* 260) which oxidizes in air to give tetradeuterioperylene along with other dehydrogenated products. We believe that these spectroscopic and chemical results demonstrate that the reaction of perylene (**2**) with sodium does in fact form a tetraanion.

The ¹H NMR bands summarized in Table I emphasize the highly aromatic diatropic character of the neutral systems; even more so, our findings point out the aromatic character of the two quadruply charged species as manifested by their diatropicity.^{11,12}

Additional studies including ¹³C and ²³Na NMR spectroscopies of these systems are under way for the purpose of gaining a deeper insight into these multiply charged species.

(11) The well-accepted correlation of Schaeffer and Schneider¹² predicts that the centers of gravity of the ¹H NMR absorptions for **3** and **4** should appear at ca. 4.2 ppm. The data shown (Table I) emphasize the enhanced diatropicity of these tetraanions.

(12) Schaeffer, T.; Schneider, W. J. *Can. J. Chem.* 1963, 41, 966-974.

Synthesis and Structure of Bis(2,4,6-tri-*tert*-butylphenyl)diphosphene: Isolation of a True "Phosphobenzene"

Masaaki Yoshifuji,* Ichiro Shima, and Naoki Inamoto*

Department of Chemistry, Faculty of Science
The University of Tokyo, Hongo, Tokyo 113, Japan

Ken Hirotsu and Taiichi Higuchi

Department of Chemistry, Faculty of Science
Osaka City University, Sumiyoshi, Osaka 558, Japan

Received January 19, 1981

In 1877 Köhler and Michaelis¹ reported the isolation of a compound which was thought to be the phosphorus analogue of azobenzene, "C₆H₅P=PC₆H₅". The phosphobenzene structure for the compound, however, was proved to be incorrect. In the early 1960's the compound was shown to have an oligomeric structure by means of molecular weight determination² and subsequently by X-ray crystallographic analysis.³

A number of reports on cyclophosphine (RP)_{*n*} (*n* = 3-6) have appeared in the literature since the structure of the phenyl derivative was elucidated.⁴ Up to date the smallest cyclophosphine is (C₂F₅P)₃.⁵ The presence of the monomeric and dimeric species

(1) Köhler, H.; Michaelis, A. *Ber. Dtsch. Chem. Ges.* 1877, 10, 807.

(2) Kuchen, W.; Grünewald, W. *Chem. Ber.* 1965, 98, 480.

(3) (a) Daly, J. J.; Maier, L. *Nature (London)* 1964, 203, 1167. (b) *Ibid.* 1965, 208, 383.

(4) As for cyclophosphines, see review articles: (a) Cowley, A. H.; Pinnell, R. P. *Top. Phosphorus Chem.* 1967, 4, 1. (b) Haiduc, I. In "The Chemistry of Inorganic Ring Systems"; Wiley-Interscience: London, 1970; Part 1, p 82. (c) Maier, L. *Org. Phosphorus Compd.* 1972, 1, 339.

(5) Cowley, A. H.; Furttsch, T. A.; Dierdorf, D. S. *J. Chem. Soc., Chem. Commun.*, 1970, 523.

(3) Müllen, K. *Helv. Chim. Acta* 1978, 61, 2307-2317.

(4) Lawler, R. G.; Ristagno, C. V. *J. Am. Chem. Soc.* 1969, 91, 1534-1535.

(5) The dianion of tetracene (18π electrons) shows ¹H NMR bands at high field.⁴

(6) "Dictionary of Hückel Molecular Orbitals (HMO) Heilbronner and Straub"; Springer Verlag: Berlin, 1966; Section 8, 16, 20.

(7) In THF-*d*₈, recorded with the aid of Bruker WH-300 pulsed FT spectrometer operating at 300.133 MHz, deuterium lock, Aspect 2000 computer, 32K memory.

(8) The reduction of **1** to **1**²⁻ was conducted by Müllen with lithium at -80 °C.³

(9) Du Pont 21-490 B single focusing instrument equipped with a commercial dual CI/EI source. Reagent gas was isobutane, source temperature 160 °C. The spectra were run by Dr. Shmuel Zitrin, Mass Spectrometry Unit, Criminal Identification Division, Police Headquarters, Jerusalem.

(10) The respective triply charged species being a radical anion could not be detected by ¹H NMR spectroscopy.